

Dynamic Oscillations of Magnetization in High Spin Magnetic Clusters

Indranil Rudra and S. Ramasesha

Solid State and Structural Chemistry Unit

Indian Institute of Science, Bangalore 560012, India

(February 1, 2008)

We have studied the time evolution of magnetization of a high spin magnetic cluster in the ground state, in the presence of a sinusoidal axial magnetic field and a static transverse field by explicitly solving time dependent schrödinger equation. We observe oscillations of magnetization in the plateau region which has all the characteristics reminiscent of the oscillation of probability of occupation of a state in a two level lattice in the presence of an oscillating electric field. The high spin of the ground state leads to a large but finite two-level pseudo-lattice. The oscillations in magnetization occur at amplitude of magnetic fields achievable in a laboratory and they also persist for a wide range of spin-dipolar interactions. Thus, our study provides a magnetic analog of the optical two-level lattice.

PACS numbers: 75.45.+j, 75.20.-g, 75.50.Xx, 75.40.Gb, 42.50.Gy

In recent years, synthesis of high nuclearity transition metal complexes in high-spin ground state has spurred interest in magnetism on a nanoscale [1]. The synthesis of Mn_{12} and Fe_8 clusters in $S=10$ ground state has led to extensive study of quantum resonant tunneling and quantum interference phenomena [2]. The quantum resonant tunneling manifests as plateaus in the magnetization *vs* magnetic field plots, with the width and location of plateaus being determined by the ramping speed of the magnetic field as well as the initial state [3]. The quantum interference phenomena observed in the Fe_8 cluster is because the paths connecting the $M_S = +10$ and $M_S = -10$ could interfere in the presence of a magnetic field, leading to an oscillation in the tunneling probabilities [4].

In these systems, the magnetic interaction between transition metal ions in neighbouring molecules is weak due to the bulky nature of the molecular complex. Thus, we can treat these molecules as isolated magnetic clusters. These clusters are characterized by exchange couplings between magnetic ions that are frustrated, leading to a high spin ground state separated from other low-lying high spin states by rather small energy gaps. These magnetic clusters also exhibit signatures of spin dipolar and other higher order spin-spin interactions allowed by the symmetry of the cluster.

The exchange Hamiltonian of the Mn_{12} system with which we shall be concerned in this letter can be written as

$$\hat{H}_{\text{exch}} = \sum_{\langle ij \rangle} J_{ij} \hat{S}_i \cdot \hat{S}_j \quad (1)$$

where the exchange interactions are given in fig.1. The model exact $S=10$ ground state as well as the low-lying excited states of this Hamiltonian have been obtained from this model by using all the spatial symmetries and

partial spin symmetry adaptation. The ground state and the first few low-lying states live in different symmetry subspaces and do not mix unless the molecular symmetry is broken [5]. Therefore, for computing properties such as quantum resonant tunneling at low-temperatures, it is sufficient to work with the different M_S manifolds within the $S = 10$ ground state. The spin-spin interactions allowed by the symmetry of the molecule leads to a spin Hamiltonian in the presence of an applied magnetic field given by

$$\begin{aligned} \hat{H}_{\text{dip}} = & D \hat{S}_{z,\text{total}}^2 + c (\hat{S}_{x,\text{total}}^4 + \hat{S}_{y,\text{total}}^4) \\ & + g_{\text{crown}} \vec{H}(t) \cdot \hat{S}_{\text{crown}} + g_{\text{core}} \vec{H}(t) \cdot \hat{S}_{\text{core}} \end{aligned} \quad (2)$$

where, for chemical reasons, the g factors for the Mn^{3+} and Mn^{4+} ions in the crown and core of the molecule (fig 1) are slightly different [6]. A time-dependent solution of the above Hamiltonian, starting from an initial state with $S = 10$ and $M_S = -10$ shows quantum resonant tunneling when the magnetic field is ramped. [3]

The energy level structure of the eigenstates of eqn. 2 are very similar to those encountered in the dynamic studies of two-level lattices. There has been considerable study of the two-level lattices in which Stark-Wannier effect and Bloch oscillations are observed [7,8]. In the presence of oscillatory electric fields, population trapping [9] and dynamic localization [10] have also been observed. The aim of the present study is to see if these effects can also be observed in the Mn_{12} magnetic cluster in the presence of an oscillatory magnetic field.

In the present work we have studied the magnetization oscillations under the influence of a strong axial *ac* magnetic field and a weak stationary transverse field, using explicit time evolution of an initial state. We start with an initial state which is the lowest eigenstate of the full exchange Hamiltonian (eqn. 1) with $S = 10$ and $M_S = -10$. In the absence of the quartic spin terms and

external transverse field, this is also the lowest energy state of the low-energy Hamiltonian (eqn. 2). We evolve this state with time explicitly, using the time-dependent schrödinger equation,

$$i\hbar \frac{\partial \psi(t)}{\partial t} = H_{\text{dip.}} \psi(t). \quad (3)$$

At each time step, we calculate the magnetization of the state. The value of D estimated from experiments is 0.56K, while c is 3×10^{-5} K [6]. The g_{crown} and g_{core} are respectively 1.96 and 2.00. In our studies, we have varied the amplitude of the time dependent magnetic field between 10 to 100 D . These fields range from 3.75 T to 37.5 T, for experimentally estimated D value in Mn_{12} , which are indeed attainable [11]. The transverse field is static and fixed at a value equal to one hundredth the amplitude of the time varying axial magnetic field.

The initial state in the time evolution is chosen to be the state with $S = 10$ and $M_S = -10$, which is the ground state, in the absence of the weak off-diagonal terms of $H_{\text{dip.}}$. The angular frequency, ω , of the axial field is varied between 10^{-1} and 10^{-3} radian $-D/\hbar$. The time evolution is carried out successively in steps $\Delta t = 0.1$ and the evolution is carried out over several periods of the applied field.

In fig. 2a, we show a plot of magnetization *vs* time, for the amplitude of the axial field, $H_0=50D$, and angular frequency $\omega = 10^{-2}$ radian $-D/\hbar$. We notice that the magnetization shows distinct plateaus and in each plateau, the magnetization oscillates a fixed number of times. The number of oscillations $N_{\text{osc.}}$ in each plateau is given by the ratio of half the energy gap between the $M_S = -10$ and the $M_S = -9$ states at the field H_0 and $\hbar\omega$, where ω corresponds to the angular frequency of the applied axial field. This kind of oscillations with similar dependence of $N_{\text{osc.}}$ has been theoretically observed in the two-level systems by Rotvig *et al* [12] and Raghavan *et al* [13]. Such oscillations have also been seen in the Bose-Einstein condensates in a double-well trap [14]. Rotvig *et al* observe this in the context of a two-band semiconductor superlattice in an external electric field, while Raghavan *et al* observe it in a single band model in the presence of an electric field. In these calculations, the probability in a given state shows the temporal oscillations that we see here for magnetization. Indeed, the different M_S states in the magnetic cluster can be viewed as forming a large but finite lattice. The transverse magnetic field couples the states at successive lattice sites, much as the transfer terms in the models of Rotvig *et al* and Raghavan *et al*. The time varying axial magnetic field corresponds to the applied oscillatory electric field of the two level systems. It is also worth noting that for the parameters that Raghavan *et al* use, the oscillations die down for larger lattices. It appears that the size of the pseudo-lattice provided by the magnetic cluster is not large enough for the oscillations to die down for the realistic model parameters we have chosen.

The oscillations in magnetic field that we observe are quite robust. We have observed these oscillations for other initial states with integral M_S values corresponding to the eigenstates of the $H_{\text{dip.}}$, with off-diagonal elements set to zero (fig. 2 b-d). The dependence of the magnetization on the amplitude of the magnetic field is shown in fig.3. We note that at higher fields, we see more oscillations in each plateau since the gap between successive M_S states widen with increasing amplitude. In fig. 4 b,d and f, we show the energies of the two low-lying eigenstates of $H_{\text{dip.}}$, corresponding to the axial field at that instance. We note that the jump in magnetization coincides with two states with different M_S values becoming degenerate. In the time between these coincidences, the magnetization shows small amplitude oscillations. Assuming that the system wavefunction evolves as $\exp(-i(\bar{E}_2 - \bar{E}_1)t/\hbar)$, where $\bar{E}_1$ and $\bar{E}_2$ are the average energies of the two states in question, the number of oscillation in a time period $t = 2\pi/\omega$ is given by $(\bar{E}_2 - \bar{E}_1)/\hbar\omega$. We note that the energy gap at the maximum amplitude in all the cases are the same and hence the number of oscillations in a plateau is inversely proportional to the frequency of the axial field. In fig. 4 a,c and e we show the dependence of the oscillation pattern on the frequency of the axial field with a fixed amplitude. We note that at decreasing frequencies, the number of oscillations in each plateau increases and the plateau structure itself vanishes with the magnetization following the magnetic field for higher frequency of the axial field.

We have also studied this system in the presence of an axial magnetic field with two different frequencies, given by $H_0\cos(\omega_1 t)\cos(\omega_2 t)$. This leads to a beat pattern involving the sum and difference of the two frequencies. In fig. 5, we show the time variation of magnetization in the presence of two different frequency magnetic fields. We note that one set of oscillations correspond to $N_{\text{osc.}} = \Delta E/2 \hbar(\omega_1 + \omega_2)$ while another set of oscillations have $N_{\text{osc.}} = \Delta E/2 \hbar(\omega_1 - \omega_2)$, where ΔE is the difference in energy between the $M_S=-10$ and $M_S=-9$ states when the axial field is equal to the amplitude.

The parameters for which we have carried out the calculations corresponds to Mn_{12} . However, systems such as Fe_8 also have high spin ground state although the D value is different. In order to study the effect of change in D value on the oscillations, we have carried out these calculations for several H_0/D values. In fig. 6, we show the results of our calculations for $H_0/D = 20$ and $H_0/D = 10$. The oscillations vanish ($H_0/D = 20$) and we have a M *vs* t behaviour that does not have much structure for this larger H_0/D value. However for lower H_0/D value, we find that the oscillations persist, but with much reduced amplitude. Thus, it is possible that in other high spin systems, these oscillations are found at different field amplitude.

To conclude, the magnetic clusters in high spin ground state behave like a finite lattice of two level systems and

in the presence of an oscillating axial field and static transverse field lead to predictable oscillations in the plateau regions of magnetization *vs* time plots. The field at which these oscillations can be observed fall in the range of 4 to 40 T with a frequency ranging from 1 to 10 MHz for the known high-spin magnetic clusters. While such behaviour is predicted in optical two level systems, their observation may be easier in high nuclearity high spin magnetic systems.

Acknowledgments

We thank Prof. Diptiman Sen for many helpful discussions, and the Council of Scientific and Industrial Research, India for their grant No. 01(1595)/99/EMR-II.

-
- [1] T. Lis, Acta Crystallogr. B **36**, 2042 (1980); C. Delfas, D. Gatteschi, L. Pardi, R. Sessoli, K. Wieghardt and D. Hanke, Inorg. Chem. **32**, 3099 (1993); A. L. Barra, D. Gatteschi, L. Pardi, A. Müller and J. Döring, J. Am. Chem. Soc. **114**, 8509 (1992).
 - [2] *Quantum Tunneling of Magnetization - QTM'94*, edited by L. Gunther and B. Barbara, NATO ASI Ser. E, Vol. 301 (Kluwer, Dordrecht, 1995); J. R. Friedman, M. P. Sarachik, J. Tejada and R. Ziolo, Phys. Rev. Lett. **76**, 3830 (1996); L. Thomas, F. Lioni, R. Ballou, D. Gatteschi, R. Sessoli and B. Barbara, Nature **383**, 145(1996) and references therein; W. Wernsdorfer and R. Sessoli, Science, **284**, 133(1999)
 - [3] Indranil Rudra, Diptiman Sen and S. Ramasesha, (Phys. Rev. B in press)
 - [4] W. Wernsdorfer and R. Sessoli, Science **284**, 133 (1999); Anupam Garg, Europhys. Lett., **22**, 205(1993) and references therein
 - [5] C. Raghu, Indranil Rudra, Diptiman Sen and S. Ramasesha, (Phys. Rev. B in press)
 - [6] A. L. Barra, D. Gatteschi and R. Sessoli, Phys. Rev. B **56**, 8192 (1997).
 - [7] G. H. Wannier, Phys. Rev. **117** 432 (1960), Rev. Mod. Phys., **34**, 645 (1962); J. E. Avron, Ann. Phys. (NY) **143**, 33 (1982) ; H. Fukuyama, R. A. Bari and H. C. Fogedby, Phys. Rev. B **8**, 5579 (1973) ; Z. Barticevic, M. Pacheco and F. Claro, Phys. Rev. B **51**, 14414 (1995); K. Gibb, M. M. Dignam, J. E. Sipe and A. P. Roth, Phys. Rev. B **48**, 8156 (1993)
 - [8] F. Bloch, Z. Phys. **52**, 555 (1928) ; M. Forst, G. Segschneider, T. Dekorsy and H. Kurz, Phys. Rev. B **61**, R10563 (2000); Jordan Kyriakidis and Daniel Loss, Phys. Rev. B **58**, 5568 (1998); Masato Morifuji and Chihiro Hamaguchi, Phys. Rev. B **58**, 12842 (1998)
 - [9] Girish S. Agarwal and W. Harshawardhan, Phys. Rev. A **50**, R4465 (1994) and references therein
 - [10] D. H. Dunlap and V. M. Kenkre, Phys. Rev. B **34**, 3625

(1986) ; Phys. Lett. A **127**, 438 (1988); Phys. Rev. B **37**, 6622 (1988)

- [11] M.D. Bird , S. Bole, I. Dixon, Y.M. Eyssa, B.J. Gao and H.J. Schneider-Muntau , Physica B **294**, 639 (2001)
- [12] Jon Rotvig, Antti-Pekka Jauho and Henrik Smith, Phys. Rev. Lett. **74**, 1831 (1995)
- [13] S. Raghavan, V. M. Kenkre, D. H. Dunlap, A. R. Bishop and M. I. Salkola, Phys. Rev. A **54**, R1781 (1996)
- [14] A. Smerzi, S. Fantoni, S. Giovanazzi and S. R. Shenoy, Phys. Rev. Lett. **79**, 4950 (1997)

Figure Captions

1. A schematic diagram of the exchange interactions between the Mn ions in the Mn_{12}Ac molecule. The interactions $J_1 = 215\text{K}$ and $J_2 = J_3 = 86\text{K}$ are antiferromagnetic, while $J_4 = -64.5\text{K}$ is ferromagnetic. The Mn^{4+} ions form the core while the Mn^{3+} ions form the crown.
- [5]
- 2.(a) Plot of evolution of magnetization with time (in units of $\hbar/D \times 10^{-1}$) for $H_0/D = 50$ and $\omega = 0.01$ radian $- D/\hbar$, initial state is $M_s = -10$. For (b), (c) and (d) $H_0/D = 100$ and $\omega = 0.01$ radian $- D/\hbar$, initial states are respectively $M_s = -9, -4$ and 4 .
3. (a) Plot of magnetization *vs* time when $H_0/D = 100$ and $\omega = 0.01$. For (b) $H_0/D = 200$ and $\omega = 0.01$. Initial state in both cases is $M_s = -10$. ω and time are in units of fig 2.
4. Magnetization *vs* time (a, c and e) and energy of $M_s = -10$ (solid line) and -9 (broken line) *vs* time (b, d and f) for (i) $\omega = 0.001$ (a, b) (ii) $\omega = 0.01$ (c, d) and (iii) $\omega = 0.1$ (e, f). $H_0/D = 100$ for all cases. ω and time are in units of fig 2.
5. Plot of magnetization *vs* time when $H_0/D = 100$, $\omega_1 = 0.01$ and $\omega_2 = 0.005$. ω and time are in units of fig 2.
6. Evolution of magnetization with time with different H_0/D values, ω is kept fixed at 0.01. $H_0/D = 100, 20$ and 10 respectively for (a), (b) and (c). ω and time are in units of fig 2.









